

HONEY BEES: BENEFICIAL ROBBERS!

PASHTE V. V¹ & SAID P. P²

¹Assistant Professor, Department of Rural Technology, Guru Ghasidas University, Bilaspur, Chhattisgarh, India

²Subject Matter Specialist, Department of Agricultural Processing and Food Engineering,
Krishi Vigyan Kendra, Bilaspur, Chhattisgarh, India

ABSTRACT

Honey bees are most important pollinators around the world and are major pollinators in tropical ecosystems. Honey bee visits the flowers to obtain their food and in return pollinate them. Bee pollination as a new agricultural production strategy has huge possibilities. Profitable crops like onion, sunflower, Apple and cucurbitaceous crops are specifically reliant upon or are benefited by honey bee pollination. Apart from the beekeeping products bee pollination benefits society by increasing food security and improving live hoods.

KEYWORDS: Honey Bee, Pollination, Foraging, Crops

INTRODUCTION

Pollination by animals is an important ecosystem service because crop plants accounting for 35% of global crop-based food production benefit from animal-mediated pollination. Flower-visiting insects, such as bees and butterflies, contribute to farmland biodiversity of habitats. They have an important functional role as pollinators of wild plants and several crop species (Klein *et al.*, 2007).

Bees (Hymenoptera: Apiforme) the primary pollinators for most of the crops requiring animal pollination (Delaplane and Mayer, 2000; Free, 1993; Klein *et al.*, 2007). In much of the world, the cornerstone of agricultural pollination is the managed honey bee (*Apis mellifera*). Honey bees, which are native to Europe and Africa but not North America, are maintained in hives which are moved into agricultural areas during crop bloom. Despite the honey bees effectiveness as a pollinator for many crops, there are risks associated with reliance on a single, managed pollinator species. Managed honey bee stocks in the USA have decreased steadily since the 1940s, and are now at less than half their original numbers (Ellis *et al.*, 2010).

Effective pollination period (EPP) has been considered (Honsho *et al.*, 2007; Sanzol and Herrero, 2001) to be an important tool to detect factors limiting fruit set. The nature and behaviour of pollinators, quality and quantity of pollen that reach to stigma, site of pollen delivery and quantity and quality of nectar are major biotic factors influencing effective pollination and fruit set in plants (Endress, 1994; Corbett, 2003). The pollen:ovule ratio is also an indicative of degree of out-crossing and mode of pollination. Low P/O ratio indicates autogamy and high xenogamy (Cruden, 2000; Jurgens *et al.*, 2002; Stefan, 2009). Pollen load, visitation frequency and pollen delivery vary with the insect species (Ivey *et al.*, 2003; Monzon *et al.*, 2004 and Stout, 2007) and in general, visitors with high pollen load and visitation frequency are considered efficient pollinators.

The Importance of Honeybees as Pollinators

Honeybees play an important role for cross-pollination, which is important for both monoecious and dioecious plant species. Without this assistance, fruit and/or seeds would not be formed for most flowering species. Many agricultural crops such as *Aeschynomene americana* L., *Ageratum conyzoides* L., *Amomum xanthioides* Wall., *Anacardium occidentale* L., *Antigonon leptopus* Hook. *Balakara baccata* Roxb., *Castanopsis acuminatissima* Rehd., *Cinnamomum kerrii* Kosten, *Coccinia grandis* CL.Voigt, *Cocos nucifera* L., *Coffea Arabica* L., *Conyza sumatrensis* Retz are pollinated by honeybees (Suwannapong *et al.*, 2011).

Honeybees of the genus *Apis* are the most studied because of their fascinating and complex lifestyle, communication systems (Nieh, 1998; Nieh and Roubik, 1995), role as keystone pollinators of native plants, pollination of agricultural crops, and the valuable hive products that they produce, such as honey, royal jelly, bee wax, bee pollen, propolis and even bee venom.

In contrast to the sole species of honey bee, there are at least 17,000 species of native, wild bees worldwide (Michener, 2007). Many of these species visit crops (Delaplane and Mayer, 2000; Klein *et al.*, 2007), and they contribute substantially to the pollination of such crops as coffee *Coffea spp.* (e.g. (Klein *et al.*, 2003), watermelon *Citrullus lanatus* (Kremen *et al.*, 2002; Winfree *et al.*, 2007), tomato *Solanum lycopersicum* (Greenleaf and Kremen, 2006a), blueberry *Vaccinium spp.* (Cane, 1997; Isaacs and Kirk, 2010), sunflower *Helianthus annuus* (Greenleaf and Kremen, 2006b) and canola *Brassica spp.* (Morandin and Winston, 2005). These native bees provide pollination thus directly benefiting crop production. In addition, they complement in a number of ways the service provided by honey bees: biologically, by enhancing the efficacy of honey bee pollination in some cases (Greenleaf and Kremen, 2006b), and economically, by insuring against pollination shortages. Having accurate estimates of this value could improve land use planning by quantifying the costs and benefits of conserving habitat for pollinators in agricultural systems.

Insect and other organisms play major role in boosting agricultural production by significantly increasing the yields of crops, vegetables, fruits and seeds through visiting flowers and helping in pollination (Pashte and Kulkarni, 2015). Self-incompatible and cross-pollinated crops require pollinating service of efficient pollinators. Self-pollinated crops also benefit from insect pollination, that increase yield up to 30% from pollinator visits and also collection of nectar or pollen and benefit farmers from pollinators' service. Lack of pollinators causes decline in fruit and seed production (Partap, 2001).

POSITION OF HONEY BEE IN ASIA

Asia has a rich diversity of honeybee species. These include *Apis cerana*, *A. dorsata*, *A. florea*, *A. laboriosa*, *A. breviligula*, *A. binghami* and *A. andreniformis* that are indigenous to the region, whereas the European honeybee, *A. mellifera* was introduced to the region and promoted for beekeeping. The indigenous honeybees make a significant contribution to the livelihoods of the rural poor and protection of the environment through a variety of products and services (Partap, 1992). There is timely need for better management of hive honeybees such as *A. cerana* and *A. mellifera* in rare pollinator areas to increase fruit production. Information on the role of honeybees in pollination leads to increased quality and yield of crops worldwide (McGreger, 1976; Crane, 1991; Free 1993; Partap and Verma, 1994; Suwannapong *et al.*, 2011).

FORAGING BEHAVIOUR

Honeybees collect the nectar and pollen of flowers as food for their colony. When performing this task, honeybees show flower constancy (Abrol); individual honeybees exclusively visit flowers of the same species as long as nectar is provided. However, honeybees are not explicitly specialized to forage on specific flower species.

They are generalists in their ability to learn the color, odor and shape of all kind of flowers (Backhaus, 1993). Honeybees have ability to learn and remember the color, shape, and fragrance of flowers that are bountiful in these nutrients, and also how to get to them. A bee can learn a new color in about half an hour (after it has made about five visits to collect a food reward), a new pattern in about half a day (after 20–30 rewarded visits), and a new route to a food source in about 3 to 4 visits. It can learn to visit different species of flowers at different locations, at different times of the day (Srinivasan, 2010).

There are differences among flowering plant species with respect to nectar and pollen production. Not all plant species possess nectaries (glands secreting nectar) or have nectar that bees can reach with their proboscis (tongue) (Partap, 1992). Nectaries can be located in various areas of the flower and some species have extrafloral nectaries that may be visited by bees. In addition, some bees may perform nectar robbing, making a small hole at the base of a flower in order to obtain the nectar. In this case, the bee does not perform any pollination service for the “robbed” plant.

Honeybees are unlikely to make many repeat visits if a plant provides little reward. A single forager will visit different flowers in the morning and, if there is sufficient attraction and reward in a particular kind of flower, she will make visits to that type of flower for most of the day, unless the plants stop producing reward or she detects forage-marking pheromones left by other bees to avoid revisiting the nectar-depleted flower.

Honeybee body are covered with abundant setae which pollen grains are attached while she forage both for nectar and pollen. They make pollen to be pellets with nectar and carry them by pollen baskets on the hind tibiae and storage in the pollen storage area (Partap, 1992). In addition to collecting nectar and pollen, foragers can collect plant gum (propolis) and water (Marcucci, 1995; Bankova *et al.*, 1983, 2000). All crops which are mostly visited and benefited by bees are reviewed in Table 1.

At the present day, there is now a general increased awareness for habitat management that may help pollinators likely to increase. Discovering potential pollinators, devising management techniques, and increasing their population for commercial exploitation will take several years. Research studies are needed in this direction to conserve honeybees and other natural pollinators, exploit their potentiality in crop pollination and allow them to develop in the pollution free environment.

CONCLUSIONS

The effectiveness of honeybees in crop pollination depends on many factors, such as the location of the hives, the attractiveness of the flowers to the bees, and the bees' behaviour in approaching the flowers. The number of colonies varies considerably according to the crop: many fruit crops require several strong hives per hectare. To ensure that a large proportion of the flowers are visited, the bees should be moved to the area as soon as blossoming begins or, preferably, just before.

A number of factors can to some extent limit the efficiency of bees as pollinators: competition from nearby

flowering plants, such as weeds; colony weaknesses brought about by the effects of pesticides, bee diseases and negligence during the pollen flow; and the effects of poor weather, which may prevent the bees from flying. Special emphasis should be placed on the danger of pesticides: while colonies of bees are in the field for pollination, toxic pesticides must not be used in the area, and in this regard, cooperation between growers and beekeepers is of the utmost importance.

REFERENCES

1. Abel, C. A., Wilson, R. L. and Luhman, R. L. (2003). Pollinating efficacy of *Osmia cornifrons* and *Osmia lignaria* sub sp. *lignaria* on three Brassicaceae species grown under field cages. *Journal of Entomological Sciences* 38(4): 545-552.
2. Ahmed, B. and Rehman, A. (2002). Population dynamics of insect foragers and their effect on seed yield of rapeseed (*Brassica campestris* L. var. toria). *Indian Bee Journal* 64 (3 & 4): 1-5.
3. Annelise de Souza Rosa, Betina Blochtein and Diego Kweco Lima (2011). Honey bee contribution to canola pollination in Southern Brazil. *Scientia Agricola* (Piracicaba) 68:255-259.
4. Bankova, V. S., Popov, S. S. and Marekov, N. L. (1983). A study on flavonoids of propolis. *Journal of Natural Products* 46: 471-474.
5. Bankova, V., Castro De, S. L. and Marcucci, M. C. (2000). Propolis: recent advances in chemistry and plant origin. *Apidologie* 31: 3-15.
6. Bisht, D. S., Naim, M. and Mehrotra, K. N. (1983). Studies on role of honey bees in rape seed production. *Proceedings of the Second International Conference Apiculture Tropical Climates* Feb 29- March 4. New Delhi, pp- 491-496.
7. Cane, J. H. (1997). Lifetime monetary value of individual pollinators: the bee *habropoda laboriosa* at rabbiteye blueberry (*Vaccinium asheireade*). *Acta Horticulturae* 446: 67-70.
8. Chand, H. and Singh, B. (1995). Effect of pollination by *Apis cerana* F. on yield of mustard, *Brassica juncea* L. *Indian Bee Journal*, 57 (1): 173-174.
9. Corbett, S. A. (2003). Nectar sugar content: estimating standing crop and secretion rate in the field. *Apidologie* 34: 1-10.
10. Crane, E. (1991). *Apis* species of tropical Asia as pollinators and some rearing methods for them. *Acta Horticulture* 288: 29-48.
11. Cruden, R. W. (2000). Pollen grains: why so many? *Plant Systematic and Evolution* 222: 143-165.
12. Delaplane, K. S. and Mayer, D. F. (2000). Crop pollination by bees. CABI publishing.
13. Delaude, A., Tasei, J. N. and Roller, M. (1978). Pollinator insects of sunflower (*Helianthus annuus* L.) in France, pollination of male sterile lines of hybrid seed production. *Proceedings of the IVth International Symposium on Pollination*, Maryland, pp.29-40.
14. Ellis, J. D., Evans, J. D. and Pettis, J. (2010). Colony losses, managed colony population decline, and colony collapse disorder in the United States. *Journal of Apicultural Research* 49 (1): 134-136.

15. Elstrom, G. W. and Maynard, D. N. (1991). Attraction of honey bees to watermelon with bee attractants. *Proceedings of Florida State Horticultural Society* 103: 130-133.
16. Endress, P. K. (1994). *Diversity and Evolutionary Biology of Tropical Flowers*. Cambridge University Press, Cambridge.
17. Free, J. B. (1993). *Insect pollination of crops*. Academic Press, London-New York.
18. Free, J. B. (1993). *Insect pollination of crops*. London: Academic Press. pp. 684.
19. Free, J.B. (1970). *Insect pollination of crops*. Academic Press, London and New York.
20. Froster, I. W., Clinch, P. G. and Jones, P. T. (1973). Observation on the pollination of Chou moellier (*Brassica napus* L.). *New Zealand Journal of Experimental Agriculture*, 1(1): 46-48.
21. Furgula, B., Noetzel, D. M. and Robinson, R. G. (1979). Observation on pollination of hybrid sunflowers. *Proceedings of the IVth International Symposium on Pollination*, Maryland, pp- 45-48.
22. Ganapathi, K. (2005). Impact of bee attractants on bee activity and yield parameters of Bt cotton. Thesis, M. Sc. [Agri.] University of Agriculture Sciences, Dharwad, Karnataka, India.
23. Goyal, N. P., Manjit, S. and Mishra, J. L. (1989a). Role of insect pollination in seed production of carrot (*Daucus carota* L.). *Indian Bee Journal* 6 (3): 89-93.
24. Greenleaf, S. S. and Kremen, C. (2006a). Wild bee species increase tomato production and respond differently to surrounding land use in northern California. *Biological Conservation* 133 (1): 81–87.
25. Greenleaf, S. S. and Kremen, C. (2006b). Wild bees enhance honey bees' pollination of hybrid sunflower. *Proceedings of the National Academy of Sciences* 103 (37):13,890-13,895.
26. Guruprasad, G. S. (2001). Maximisation of niger productivity through enhancement of bee pollination. Thesis, M. Sc. [Agri.] University of Agriculture Sciences, Dharwad, Karnataka, India.
27. Harichand, and Singh, R. (1995). Effect of pollination by *Apis cerana* Talor. on yield of mustard, (*Brassica juncea* Linn). *Indian Bee Journal* 57: 173-174.
28. Honsho, C., Songpol, S., Takuya, T., Kensuke, Y. and Keizo, Y. (2007). Effective pollination period in durian (*Durio zibethinus* Murr.) and the factors regulating it. *Scientia Horticulturae* 111:193-196.
29. Isaacs, R. and Kirk, A. K. (2010). Pollination services provided to small and large high bush blueberry fields by wild and managed bees. *Journal of Applied Ecology* 47 (4) 841-849.
30. Ivey, C. T., Martinez, P. and Wyatt, R. (2003). Variation in pollinator effectiveness in swamp milkweed *Asclepias incarnate* (Apocynaceae). *American Journal of Botany* 90: 214-225.
31. Jurgens, A., Witt, T. and Gottsberger, G. (2002). Pollen grain numbers, ovule numbers and pollen ovule ratios in Caryophylloideae: correlation with breeding system, pollination, life from, style number, and sexual system. *Sexual Plant Reproduction* 14: 279-289.
32. Kermer, J. C. (1945). Influence of honeybees habits on radish seed yield. *Michigan Agricultural Experimental*

Station Quarterly Bulletin 27: 413-428.

33. Khan, B. M. and Chaudhary, M. I. (1988). Comparative assessment of honeybees and other insects with self-pollination of sarson in Peshawar. *Pakistan Journal of Forestry* 38 (4): 231-237.
34. Klein, A. M., Vaissiere, B. E., Cane, J. H., Steffan-Dewenter, I., Cunnigham, S. A., Kremen, C. and Tscharntke, T. (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B London: Biological Sciences* 274: 303-313.
35. Klein, A. M., Steffan-Dewenter, I. and Tscharntke, T. (2003). Pollination of *coffea canephora* in relation to local and regional agroforestry management. *Journal of Applied Ecology* 40 (5), 837-845.
36. Kremen, C., Williams, N. M. and Thorp, R. W. (2002). Crop pollination from native bees at risk from agricultural intensification. *Proceedings of the National Academy of Sciences* 99 (26): 16812.
37. Kulkarni, S. N. and Dhanorkar, B. K. (1998). Effect of *Apis cerana indica* on niger seed production in Marathawada region. Paper presented at *FAO Workshop on Sustainable Bee Keeping Development*, 1-5 August, Dharwad.
38. Kumar, M., Chand, H., Singh, R. and Ali, M. (2002). Effect of different modes of honeybee pollination on oil content in seeds of sunflower (*Helianthus annus* L.). *Journal of Entomological Research* 26 (3): 219-221.
39. Langdrige, D. F. and Goodman, R. D. (1975). A study in pollination of oil seed rape. *Australian Journal of Experiments in Agriculture and Animal Husbandry*. 22: 124-126.
40. Langridge, D. F. (1969). The role of honeybees in pollination of apples. *Apiacta* 3.
41. Lerin, J. (1982). Effects of insect pollination on rape in a cage experiment. *Agronomie* 2 (3): 249-255.
42. Lingappa, S., Viraktmath, S. A., Vastrad, A. S. and William, R. (1999). Utilization of honey bee *Apis cerana indica* F. for pollination of water melon and safflower. *Proceedings of Apimondia*, Pp.235.
43. Marcucci, M. C. (1995). Propolis: chemical composition, biological properties and therapeutic activity. *Apidologie* 26: 83-99.
44. McGregor, S. E. (1976). Insect pollination of cultivated crop plant. *Agricultural Handbook*. Washington D.C.: USDA-ARS. New York. pp. 496.
45. Michener, C. D. (2007). *The bees of the world*, Second ed. Johns Hopkins University Press, Baltimore, Pp-992.
46. Mishra, R. C, Jitender Kumar and Gupta, J. K. (1988). The effect of mode of pollination on yield and oil potential of *Brassica campestris* L. Var. Sarson with observation on insect pollinators. *Journal of Apicultural Research* 27 (3): 186-189.
47. Mohammad, S. M., Shazia, R., Shahid, N. and Ghulam, S. (2011). Comparative performance of honeybees (*Apis mellifera* L.) and blow flies (*Phormia terronovae*) in onion (*Allium cepa* L.) seed setting. *Journal of Agriculture Research* 49(1):49-56.
48. Monzon, V. H., Bosch, J. and Retana, J. (2004). Foraging behaviour and pollinating effectiveness of *Osmia cornuta* (Hymenoptera: Megachilidae) and *Apis mellifera* (Hymenoptera: Apidae) on "Comice" pear. *Apidologae*

35: 575-585.

49. Morandin, L. A. and Winston, M. L. (2005). Wild bee abundance and seed production in conventional, organic, and genetically modified canola. *Ecological Applications* 15(3): 871–881.
50. Murasing, S. (2000). Maximization of niger productivity through enhancement of bee pollination. Thesis, M. Sc [Agri.] University of Agriculture Sciences, Dharwad, Karnataka, India.
51. Nieh, J. C. (1998). The role of scent beacon in the communication of food location by the stingless bee, *Melipona panamica*. *Behaviour Ecology Sociobiology* 43: 47-58.
52. Nieh, J. C. and Roubik, D. W. (1998). Potential mechanisms for the communication of height and distance by a stingless bee, *Melipona panamica*. *Behaviour Ecology Sociobiology* 43: 387-399.
53. Partap, T. (1992). *Honey plant sources in mountain areas*. Honeybee in Mountain in Agriculture. New Delhi: Oxford and IBH. pp. 91-112.
54. Partap, T. (2001). Mountain agriculture, marginal land and sustainable livelihoods: Challenges and opportunities. *International Symposium on Mountain Agriculture in HKH Region*, 21-24 May 2001, ICIMOD, Kathmandu, Nepal.
55. Partap, U. and Verma, L. R. (1994). Pollination of radish by *Apis cerana*. *Journal of Apicultural Research* 33: 237-241.
56. Pashte, V. V. and Kulkarni, S. R. (2015). Role of Pollinators in Qualitative Fruit Crop Production: A Review. *Trends in Biosciences* 8(15): 3743-3749.
57. Pashte V. V. and Shylesha A. N. (2013a). Pollinators diversity and their abundance on sesamum. *Ind J Entomol* 75: 260-261.
58. Pashte V. V. and Shylesha A. N. (2013b). Effect of number of honey bee visits on yield of sesamum. *Bioinfolet* 10: 1591-1592.
59. Pashte V. V. and Shylesha A. N. (2013c). Effectiveness of attractants and scents in enticement of *Apis dorsata*, *Apis florea* and non-apis bees on sesamum (*Sesamum indicum*). *Bioinfolet* 10(4C): 1593-1596.
60. Pashte V. V. and Shylesha A. N. (2013d). Pollen and nectar foraging activity of honey bees in sesamum. *Indian Journal of Entomology* 75(2): 124-126.
61. Patil, B. S. (1999) Role of bee attractant in enhancing productivity and quality of sesamum. Thesis, M. Sc [Agri.] University of Agriculture Sciences, Dharwad, Karnataka, India.
62. Prasad, D., Hameed, S. F. Singh, R., Yazdani, S. S. and Singh, B. (1989). Effect of bee pollination on the quantity and quality of rai crop (*Brassica juncea* Coss.). *Indian Bee Journal* 51(2): 45-47.
63. Priti, R. C., Mishra, and Sihag, R. C. (2001) Role of insect pollination in seed production of radish (*Raphanus sativus* L.). *Seed Research* 29(2): 231-234.
64. Rakesh Kumar, and Lenin, L. K. (2000). Insect pollinators and effects of cross pollination on yield attributes of sesame (*Sesame indicum*). *Indian Bee Journal* 62 (1-2): 75-80.

65. Rakeshkumar, O. P., Chaudhary, Lenin, L. K. (1994). Studies on the foraging behavior of honeybees and their role as pollinators of sunflower (*Helianthus annuus* L.). *Indian Bee Journal* 56: 207-210.
66. Ramachandran, T. K. and Menon, M. P. (1979). Pollination mechanism and inbreeding depression in niger. *Madras Agril. Journal* 66: 449-454.
67. Sanzol, J. and Herrero, M. (2001). The “effective pollination period” in fruit trees. *Scientia Horticulturae* 90: 1-17.
68. Sattagi, H. N., Kambrekar, D. N., Gundannavar, K. P. and Kulkarni, K. A. (2004). Effect of bee pollination on the yield parameters of niger, *Karnataka Journal of Agriculture Sciences* 17(4): 831-832.
69. Sattagi, H. N., Rajashekhar, D. W. and Kulkarni, K. A. (2001a). Effect of attractants in enhancing the productivity of watermelon. Paper presented at *National Symposium on environment and Evolutionary Biology*, 1-3 March, Dharwad, 2001, Pp.24.
70. Sattagi, H. N., Rajashekhar, D. W., Kulkarni, K. A. (2001b). Effect of bee pollination on niger seed production. Paper presented at *National Symposium on Environment and Evolutionary Biology*, 1-3 March, Dharwad, 2001, Pp.37.
71. Sihag, R. C. (1986). Insect pollination increases seed production in cruciferous and umbelliferous crops. *Journal of Apicultural Research*, 25: 121-126.
72. Singh, B. B., Chand, H., Singh, S. P. N. and Kumar, M. (2011). Foraging profiles of different species of honey bees on toria (*Brassica campestris*). *Journal of Entomological Research* 35(3): 251-254.
73. Singh, A. K. and Chamotre, V. K. (1992). Seed setting under different system of pollination in radish. *Vegetable Science*, 19 (2): 217-220.
74. Singh, J., Mishra, R. C. and Agarwal, O. P. (2002). Pollinating efficiency of *Apis* species in hybrid seed production of *Brassica napus* L. *Indian Bee Journal* 64 (4): 62-63.
75. Solange, A. B., Dos S., Roselino, A. C. and Luci, R. B. (2008). Pollination of cucumber, *cucumis sativus* L. (Cucurbitales: Cucurbitaceae), by the stingless bees *Scaptotrigona* aff. *depilis* moure and *Nannotrigona testaceicornis* lepeletier (Hymenoptera: Meliponini) in greenhouses. *Neotropical entomology* 37(5): 506-512.
76. Stefan, G. M. and Walter, D. (2009). Pollination mode and life form strongly affect the relation between mating system and pollen to ovule ratios. *New Phytologist* 183: 470-479.
77. Stout, J. C. (2007). Pollination of invasive *Rhododendron ponicum* (Ericaceae) in Ireland. *Apidologie* 38:198–206.
78. Suhail, A., Zain-ul-abdin, Iqbal, J., Iqbal, M., Weseem, U., Shahid, R. N. and Ikram-ul-haq (2001). Insecticidal mortality and pollination role of honeybee (*Apis mellifera* L.) on cucumber (*Cucumis sativus* L.) crop. *International journal of agriculture & biology* 3(4):501-502.
79. Sunder Rajurs, S. (2001). Pollination potentiality of honey bees in niger production. Thesis, M. Sc [Agri.] University of Agriculture Sciences, Bangalore, Karnataka, India.
80. Suwannapong, G., Yemor, T., Boonpakdee, C. and Benbow, M. E. (2011). *Nosema ceranae*, a new parasite in

- Thai honeybees. *Journal of Invertebrate Pathology* 106:236-241.
81. Swaminathan, R. and Bharadwaj, S. C. (1998). Role of insect pollinators in seed setting of sunflower (*Helianthus annuus* L.) at Udaipur, Rajasthan, India. *Indian Bee Journal* 60: 155-156.
 82. Uma, and Verma, L. R. (1994). Pollination of radish by *Apis cerana*. International Centre for Integrated Mountain Development (ICIMOD), Kathmandu, Nepal.
 83. Vaishampayan, S. and Sinha, S. N. (2000). Effect of caging on bee behaviour and pollination efficiency of *Apis mellifera* on seed production of hybrid sunflower. *Agriculture Science Digest* 20 (2): 81-83.
 84. Verma, L. R. and Partap, U. (1994). Foraging behaviour of *Apis cerana indica* on cauliflower and cabbage and its impact on seed production, *Journal of Apicultural Research* 33 (4): 231-236.
 85. Verma, S. K. and Phogat, K. P. S. (1994). Impact of pollination by honey bees (*Apis cerana*) on yield of radish under valley condition of Himalayan hills. *Indian Bee Journal*, 45: 183-186.
 86. Verma, S. K. and Joshi, N. K. (1983). Studies on the role of honeybees in the pollination of cauliflower (*Brassica oleracea* L. var. Botrytis). *Indian Bee Journal*, 45 (3): 45-55.
 87. Viraktmath, S. A. and Anagoudar, J. A. (2002). Influence of bee attractants in enhancing pollination and yield parameters in *Cucumis sativa* L. *Indian Bee Journal* 64 (1&2): 23-27.
 88. Viraktmath, S. A. and Patil, R. K. (2002). Effect of bee pollination and change in crop environment on seed yield of sunflower. *Indian Bee Journal* 64 (1&2): 35-42.
 89. William, R. D. (2003). Exploration of *Apis cerana* F. for enhancing the productivity of sunflower safflower and watermelon. Thesis, M. Sc [Agri.] University of Agriculture Sciences, Dharwad, Karnataka, India.
 90. Wilson, K. L and Collison, V. L. (1988). Field cage study of the effects of four honey bee strains and hand pollination on the seed of a wild sunflower. *Seed Science and Technology* 16: 471-475.
 91. Winfree, R., Williams, N. M., Dushoff, J., Kremen, C. (2007). Native bees provide insurance against ongoing honey bee losses. *Ecology Letters* 10(11):1105-1113.
 92. Zdzisaw Wilkaniec, Karol giejdasz and Grzegorz Proszynski (2004). Effect of pollination of onion seeds under isolation by the red mason bee (*Osmia rufa* L.) (*Apoidea, megachilidae*) on the setting and quality of obtained seeds. *Journal of apicultural science* 48(2):35-41.

APPENDICES

Table 1: Crops Mostly Visited and Benefitted by Bees

Radish	Kermer (1945), Verma and Poghat (1994), Uma and Verma (1994), Priti <i>et al.</i> (2001), Kapila <i>et al.</i> (2002)
Mustard	Annelise de Souza Rosa <i>et al.</i> (2011), Singh <i>et al.</i> (2011), Landridge and Goodman (1975), Froster <i>et al.</i> (1973), Abel <i>et al.</i> (2003), Lerin (1982), Bisht <i>et al.</i> (1983), Prasad <i>et al.</i> (1989), Singh and Chamotre (1992), Chand and Singh (1995), Verma and Joshi (1983), Sihag (1986), Khan and Chaudhary (1988), Mishra <i>et al.</i> (1988), Harichand and Singh (1995), Singh <i>et al.</i> (2002), Murasing (2000), Ahmed and Rehman (2002)
Safflower	William (2003), Lingappa <i>et al.</i> , (1999)

Table 1: Contd.,	
Sunflower	Free (1970), Furgula <i>et al.</i> (1979), Rakeshkumar <i>et al.</i> (1994), Swaminathan and Bharadwaj (1998), Vaishampayan and Sinha (2000). (Kumar <i>et al.</i> 2002), Viraktmath and Patil (2002)
Cotton	Ganapathi, 2005
Niger	Guruprasad (2001), Ramachandran and Menon (1979), Kulkarni and Dhanorkar (1998), Sattagi <i>et al.</i> (2004), Sunder rajurs (2001), Sattagi <i>et al.</i> (2001b),
Carrot	Goyal <i>et al.</i> (1989), Kumar <i>et al.</i> (1989)
Watermelon	Elstrom and Maynard (1991), Sattagi <i>et al.</i> (2001a)
Cauliflower	Verma and Partap (1994)
Sesamum	Patil (1999), Rakeshkumar and Lenin (2000), Pashte and Shylesha (2013a), Pashte and Shylesha (2013b), Pashte and Shylesha (2013c), Pashte and Shylesha (2013d),
Cucumber	Suhail <i>et al.</i> (2001), Solange <i>et al.</i> (2008)
Onion	Mohammad <i>et al.</i> (2011), Zdzisaw Wilkaniec <i>et al.</i> (2004)
Apple	Langridge (1969)